

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Sulfur isotope analysis of microcrystalline iron sulfides using SIMS imaging: Extracting local paleo-environmental information from modern and ancient sediments.

Permalink

<https://escholarship.org/uc/item/7sj7b8ms>

Authors

Bryant, Roger N
Jones, Clive
Raven, Morgan R
[et al.](#)

Publication Date

2018-12-18

DOI

10.1002/rcm.8375

Peer reviewed

1Sulfur isotope analysis of microcrystalline iron sulfides using SIMS imaging: Extracting 2local paleo-environmental information from modern and ancient sediments

3Roger N. Bryant^{1,*}, Clive Jones¹, Morgan R. Raven^{1,2}, Maya L. Gomes³, William M.

4Berelson⁴, Alexander S. Bradley¹, David A. Fike¹

5¹Department of Earth & Planetary Sciences, Washington University in Saint Louis, St. Louis,
6MO 63130, USA.

7²Department of Earth Science, University of California, Santa Barbara, CA 93106-9630,
8USA.

9³Department of Earth & Planetary Sciences, Johns Hopkins University, Baltimore, MD
1021218, USA.

11⁴Department of Earth Sciences, University of Southern California, Los Angeles, CA 90089-
120740, USA.

13*Correspondence to: mbryant@wustl.edu.

14

15**RATIONALE:** Sulfur isotope ratio measurements of bulk sulfide from marine sediments
16have often been used to reconstruct environmental conditions associated with their formation.
17In-situ microscale spot analyses by secondary ion mass spectrometry (SIMS) and laser
18ablation multiple-collector inductively coupled plasma mass spectrometry (LA-MC-ICP-MS)
19have been utilized for the same purpose. However, these techniques are not often suitable for
20small ($\leq 10 \mu\text{m}$) grains or for detecting intra-grain variability.

21**METHODS:** Here, we present a method for the physical extraction (using lithium
22polytungstate heavy liquid), and subsequent sulfur isotope analysis (using SIMS; CAMECA
23IMS 7f-GEO) of microcrystalline iron sulfides. SIMS sulfur isotope ratio measurements were
24made via Cs^+ bombardment of raster squares with sides of 20-130 μm , using an electron
25multiplier (EM) detector to collect counts of $^{32}\text{S}^-$ and $^{34}\text{S}^-$ for each pixel (128x128 pixel grids)
26for between 20 and 960 cycles.

27**RESULTS:** The extraction procedure did not discernibly alter pyrite grain-size distributions.
28Apparent inter-grain variability in $^{34}\text{S}/^{32}\text{S}$ in 1-4 μm -sized pyrite and marcasite fragments
29from isotopically homogeneous hydrothermal crystals was $\sim \pm 2\text{‰}$ (1σ), comparable to the
30standard error of the mean for individual measurements ($\leq \pm 2\text{‰}$, 1σ). In contrast, grain-
31specific $^{34}\text{S}/^{32}\text{S}$ in modern and ancient sedimentary pyrites and marcasites can have inter- and
32intra-grain variability $> 60\text{‰}$. Distributions of intra-sample isotopic variability are consistent
33with bulk $^{34}\text{S}/^{32}\text{S}$ values.

34**CONCLUSIONS:** SIMS analyses of isolated iron sulfide grains yielded distributions that are
35isotopically representative of bulk $^{34}\text{S}/^{32}\text{S}$ values. Populations of iron sulfide grains from
36sedimentary samples record the evolution of the S-isotopic composition of pore water sulfide
37in their S-isotopic compositions. These data allow past local environmental conditions to be
38inferred.

39 Iron sulfide minerals, particularly pyrite (FeS_2), represent a substantial geologic reservoir of
40 sulfur. Pyrite is a key constituent of many iron sulfide ore deposits,¹ a common accessory
41 phase in an array of igneous and metamorphic rocks,^{2,3} and a nearly ubiquitous mineral in
42 marine sedimentary rocks of all ages.⁴ Sedimentary pyrites have diverse morphologies,
43 crystal sizes, and S-isotope compositions, and these characteristics have proven to be
44 invaluable archives for reconstructing local environmental conditions as well as global-scale
45 changes in biogeochemistry.⁵⁻⁸ The S-isotopic composition of pyrite is expressed here in the
46 standard delta notation (in units of per mil, ‰) relative to the Vienna Canyon Diablo Troilite
47 (VCDT) reference standard for sulfur,⁹

$$48 \quad \delta^{34}\text{S}_{\text{sample}} = \left(\frac{R_{\text{sample}}}{R_{\text{VCDT}}} - 1 \right) \quad (1)$$

49 where R represents $^{34}\text{S}/^{32}\text{S}$ ratios.

50 Pyrite $\delta^{34}\text{S}$ values are commonly obtained by bulk extraction of chromium-reducible
51 sulfur (CRS), using Cr^{2+} to reduce and volatilize FeS_2 to H_2S gas, which can subsequently be
52 trapped as zinc or silver sulfide.^{10,11} Isotope ratios for the isolated CRS can be measured by
53 combustion elemental analysis isotope ratio mass spectrometry (EA-IR-MS),¹²⁻¹⁷ or by
54 solution multiple-collector inductively coupled plasma mass spectrometry (MC-ICP-MS).¹⁸⁻²¹
55 Such approaches yield an integrated $\delta^{34}\text{S}$ value for the CRS pool and are thus blind to patterns
56 of isotopic variability within the pyrite pool. In addition, the CRS pool can potentially also
57 include marcasite (FeS_2), elemental S, organic polysulfides, and other metal sulfides in
58 addition to multiple morphologies or generations of pyrite.^{8,10,11}

59 The degree of isotopic variability within the pyrite pool is hypothesized to reflect
60 depositional conditions and may record critical information about the location(s) and
61 condition(s) of pyrite formation.^{8,22} For example, if pyrite in a sample is isotopically
62 homogeneous, this would suggest that all grains formed from the same fluid. In contrast, a
63 bimodal pyrite $\delta^{34}\text{S}$ distribution might indicate two distinct sulfide sources, e.g., from
64 successive exposure to distinct sulfide-bearing fluids. Pyrites from a single source may also
65 have a range of $\delta^{34}\text{S}$ values, representing their continuous formation in the presence of an
66 evolving sulfide reservoir.^{23,24} In diffusively limited environments like marine sediments, pore
67 water sulfate can be drawn down by microbial sulfate reduction (MSR). Sulfate reducers have
68 a strong preference for the lighter isotopes of S in sulfate,^{25,26} which leads to increasingly ^{34}S -
69 enriched sulfate and sulfide with depth in the sediment.²⁷ As a result, later-formed pyrite
70 crystals or the outer layers of large pyrite crystals may be more ^{34}S -enriched than pyrite that
71 formed earlier.²³ These distinct scenarios (unimodal distribution, bimodal distribution, or
72 evolved source) would not be distinguishable using bulk $\delta^{34}\text{S}$ values. Therefore, there is great
73 potential for a grain-specific method for pyrite S-isotope measurements to enhance our
74 understanding of pyrite $\delta^{34}\text{S}$ records, providing a new dimension of information to inform our
75 interpretations of this powerful archive.

76 Progress has been made using spot analyses by SIMS,^{1,23,28-38} or LA-MC-ICP-
77 MS,^{20,39,40} to make spatially resolved $\delta^{34}\text{S}$ measurements. Some have already attributed

78detectable intra-sample pyrite $\delta^{34}\text{S}$ variability to temporal changes in the S-isotopic
79composition of the fluids from which the pyrites precipitated.^{24,28} The method described here,
80scanning ion imaging by SIMS, is designed to address several of the limitations of currently
81available methods for micro-scale analysis of $\delta^{34}\text{S}$ in sedimentary pyrite. First, by rastering
82over grains, scanning ion imaging can generate a continuous record of isotope variations,²⁹
83one that can be interrogated at variable spatial resolution after data collection. Previously,
84most SIMS studies of pyrites relied on analyses of fairly large ($\geq 10\ \mu\text{m}$ -diameter) spots
85within grains.^{1,23,28–38} This precludes analyses of many sedimentary pyrites (i.e., those with
86diameters $< 10\ \mu\text{m}$). Moreover, while spot analyses can be sufficient to determine the
87presence of inter-grain isotopic variability on larger grains, they are not able to discern intra-
88grain variability,^{31–37} except for the case of unusually large (diameters $> \sim 100\ \mu\text{m}$) pyrites.²³

89 Additionally, by reducing the primary beam current, we achieve the spatial resolution
90necessary to measure inter- and intra-grain $\delta^{34}\text{S}$ variability in microcrystalline pyrites. In
91comparison, spot analyses are not ideal for many micro-sized pyrites or necessarily even for
92larger pyrites that are composed of many small parts (e.g., framboids). Relatively high
93primary beam currents (e.g., $\geq 1\ \text{nA}$) inherently limit the three-dimensional resolution of
94isotopic measurements.^{28,38} As scanning ion imaging requires the use of very low primary
95beam currents (e.g., $\leq 20\ \text{pA}$) to prevent the saturation of the electron multiplier detector,²⁹
96the associated low sputter rates and small diameter of the focused primary beam ($\leq 1\ \mu\text{m}$)
97result in excellent three-dimensional resolution.²⁹

98 Finally, we introduce a physical extraction procedure to enable pyrite to be
99concentrated for optimized SIMS analyses for samples where pyrite is a trace phase. As
100pyrite is not a high abundance phase in most sedimentary rocks, and grains are often broadly
101disseminated,⁴¹ this limits the number of grains likely to be present at the polished surface of
102a 1-inch round thin section. Additionally, when grains are close to the edge of the sample
103holder, this adversely affects precision.⁴² Therefore, the physical extraction of pyrites from
104sedimentary samples is in many cases a necessary precondition for efficient SIMS analysis on
105a sufficient number of grains to characterize the population. Most early physical extraction
106procedures for pyrite used magnetic separation, although because pyrite is only paramagnetic,
107these approaches are inefficient.⁴³ Alternative separation procedures utilize heavy liquids,^{44–48}
108as these represent an opportunity to separate dense phases like pyrite ($5.01\ \text{g/cm}^3$) from less
109dense, but common insoluble minerals like quartz ($2.65\ \text{g/cm}^3$) and clays ($2\text{--}2.7\ \text{g/cm}^3$). For
110this study, lithium polytungstate (LST; working density of $2.85\ \text{g/cm}^3$) is preferred due to its
111low toxicity and high recyclability.

112 In the following, we present the details of our SIMS scanning ion imaging-based
113sulfur isotope analysis procedure. We first test the validity of the method on micron-scale
114fragments of cm-scale isotopically homogeneous hydrothermal pyrite and marcasite crystals.
115By comparing SIMS results to bulk $^{34}\text{S}/^{32}\text{S}$ ratio data obtained via EA-IR-MS, we quantify
116overall analytical biases associated with the method and determine its general limitations in
117order to provide a robust platform for future grain-specific SIMS analyses of modern and
118ancient microcrystalline iron sulfides. By generating continuous $^{34}\text{S}^-$ and $^{32}\text{S}^-$ ion maps for

119inter/intra-grain variability and probing variation in hydrothermal crystals, we increase our
120confidence that the observed variation in sedimentary samples is environmentally meaningful
121and not an analytical artefact. Lastly, we apply the method to a set of extensively studied
122sediment samples that are thought to represent distinct styles and histories of pyritization,
123demonstrating its applicability and value to investigations of modern and ancient marine
124sediments.

125Materials & Methods

126Samples

127For use in density separation procedure tests, and as a S-isotopic standard for SIMS
128experiments, we obtained a single large (~2cm-diameter) euhedral pyrite crystal from Ward's
129Science (Rochester, NY), sourced from the San Jose de Huanzala Mine, Peru. The pyrite at
130this locality is of hydrothermal origin, and has been shown in previous studies to be
131stoichiometric FeS₂.⁴⁹ For use as an additional S-isotopic standard for SIMS experiments, we
132obtained a single large (~1cm-diameter) euhedral marcasite crystal from Ward's Science,
133sourced from the Jiří open-pit lignite mine in the Czech Republic. For use in further SIMS S-
134isotope experiments, we selected a modern sediment sample from Santa Barbara Basin,⁵⁰ and
135two mid-Cretaceous, Cenomanian-Turonian Ocean Anoxia Event (OAE 2) shale samples
136from the Cismon section in Italy,⁵¹ and the Demerara Rise.⁵²

137Extraction of microcrystalline pyrite from geologic samples

138A ~0.5g fragment of the hydrothermal pyrite crystal was ground to a fine powder by mortar
139and pestle, and homogenized. The resulting grain size distribution of the powdered crystal
140('pre'; Table S1) was characterized using a combination of optical microscopy and image
141processing in ImageJ.⁵³ This involved suspending ~0.001 g of powder in ethanol and
142dispersing the sample on a glass microscope slide, then capturing 20 focused images of
143unique parts of the slide through a 40× optical objective. These images were overlain by
14420×20 μm grids, and a random number generator was used to select 5 grid squares to analyze
145per image. In ImageJ, images of grid squares were scale-calibrated, converted to grayscale,
146and a bandpass filter (filtering large structures down to 10 pixels) and threshold were applied
147to highlight grains. Overlapping/touching grains, or grains with a circularity of <0.9 were
148removed, to approximate the near-sphericity of natural pyrite grains. Grain areas were
149calculated for the remaining particles using the 'Analyze Particles' function in ImageJ, and
150grain sizes were estimated by assuming that each grain was perfectly circular in cross-section.

151 A separation procedure was then carried out on subsamples of the powdered crystal.
152Three 0.01 g aliquots of powder were added to 45 mL of LST in three 50 mL centrifuge
153tubes. These tubes were mixed for a minute using a vortex mixer at the highest speed, then
154placed in an ultrasonic bath (35 kHz) for 15 minutes. Tubes were then spun in a centrifuge for
15538 minutes at 3000 rpm, as these conditions were estimated (see equation 2) to allow ≥0.5 μm
156diameter spherical pyrite fragments to settle in LST. Particle settling time (t_s) was estimated
157using Stokes' Law^{54,55}:

158
$$t_s = \frac{9 \mu h}{2(\rho_p - \rho_l) R_p^2 a} \quad (2)$$

159 where μ is the dynamic viscosity of the fluid (0.011 Pa·s for LST), h is the height of the liquid
160 in the centrifuge tube (10 cm in our setup), ρ_p and ρ_l are the mass densities of the particles and
161 the fluid (5.01 and 2.85 g/cm³ for pyrite and LST), R_p^2 is the radius of the settling particle
162 squared, and a is the centrifugal acceleration (in m/s²), given by:

163
$$a = \omega^2 r \quad (3)$$

164 where r is the centrifuge radius (18 cm in our setup), and ω is angular velocity (in radians/s),
165 given by:

166
$$\omega = 2 \pi f \quad (4)$$

167 where f is the rotational frequency (50/s in our setup). It should be noted that the settling time
168 calculated using equation 2 is likely an underestimate due to hindering effects such as
169 particle-particle and particle-container interactions. After centrifugation, settled fragments
170 were removed from the tubes using a plastic micro-pipette, placed in new 50 mL centrifuge
171 tubes, rinsed and spun down (5 minutes at 2000 rpm) five times in deionized water. This
172 process was repeated three times sequentially (Table S1, 'a-c') for each 0.01 g portion of
173 powder. In order to more closely approximate insoluble residues from marine sediments, the
174 extraction procedure was also repeated for a 1:99 mixture of pyrite and ~300 μ m-sized quartz
175 grains (0.01 g pyrite, 0.99 g quartz), and a 1:99 mixture of pyrite and ~50 μ m-sized quartz
176 grains. Each extract was dried and weighed, and grain size distributions were calculated for
177 the first sequential extracts for the pure pyrite ('post'), pyrite with large quartz grains
178 ('postQz'), and pyrite with small quartz grains ('postSmQz').

179 A shatter box was employed to powder the Cretaceous-age shale samples, for no
180 longer than one minute. There was no obvious sign of sulfide mineral fragmentation after this
181 procedure. For the Santa Barbara Basin sediment sample, one gram of dried sediment was
182 powdered gently in a mortar and pestle. For the Cretaceous shales and modern Santa Barbara
183 Basin sediment, carbonate minerals were removed by three sequential 10-minute treatments
184 with 6M hydrochloric acid, before the insoluble residue was rinsed five times with deionized
185 water and dried in an oven at 60°C for 24 hours. The insoluble residues were then powdered
186 by mortar and pestle prior to performing a single density separation as described above on
187 0.25 g aliquots of each.

188 **Mounting of samples**

189 Dried iron sulfide samples were carefully transferred onto the surface of the base of a 1-inch
190 round acrylic mould that had been coated with a release agent, isooctane. After adding iron
191 sulfide samples and powdered hydrothermal pyrite and marcasite for use as internal S-
192 isotopic standards (kept separate using a Parafilm M grid), the upper half of the mould was

193 then attached and filled with epoxy (2:1:13.63 ratio mixture of 1-(2-aminoethyl) piperazine;
194 1,8-diamino-p-menthane; and Araldite 506 epoxy resin). After degassing in a vacuum oven
195 for 10 minutes, and removing any remaining bubbles with a 21G needle, the epoxy was cured
196 for 72 h at 60°C in an oven. The epoxy was then removed from the mould and sequentially
197 polished with a 6 µm polishing pad, 3 µm diamond paste, and 1 µm diamond paste, in order
198 to expose the standards and sample, and minimize surface topography and roughness. After
199 using Raman microprobe analysis (1 mW laser power and 50x objective)⁵⁶ and optical
200 microscopy (50x objective, plane-polarized light) to confirm the presence and mineralogy of
201 analyte at the surface of the polished epoxy pucks, the pucks were coated with ~50 nm thick
202 Au to ensure conductivity for SIMS analysis.

203 **Sample imaging**

204 For iron sulfide extracts from the Cismon and Demerara shale samples, ~20 µg aliquots were
205 mounted on carbon tape, coated with 5 nm Au by physical vapor deposition using a Kurt J.
206 Lesker PVD 75 (Jefferson Hills, PA, USA), before representative mineral textures were
207 imaged by Scanning Electron Microscopy (SEM) using a JEOL JSM-7100 LVF Field
208 Emission SEM (Tokyo, Japan).

209 **Bulk sulfur isotope analyses**

210 To determine $\delta^{34}\text{S}$ values for the hydrothermal pyrite and marcasite crystals, and to assess
211 their degrees of isotopic homogeneity, three fragments of each crystal were randomly
212 selected, and powdered in an agate mortar. Small (~125 µg) aliquots of powdered FeS₂ were
213 loaded into tin capsules with 1-2 mg V₂O₅, combusted in a Costech ECS 4010 Elemental
214 Analyzer (Valencia, CA, USA); ³⁴S/³²S ratios were then measured in a Thermo Scientific
215 Delta V Plus isotope ratio mass spectrometer (Waltham, MA, USA), and corrected to VCDT
216 by bracketing analyses of in-house VCDT-calibrated ZnS, BaS and BaSO₄ standards. For the
217 fragments of pyrite and marcasite, the average $\delta^{34}\text{S}$ values measured by EA-IR-MS were -1.0
218 $\pm 0.1\text{‰}$ (1σ , $n=3$) and $4.6 \pm 0.2\text{‰}$ (1σ , $n=3$), respectively. The same analytical procedure was
219 used to generate 'bulk' $\delta^{34}\text{S}$ data for the pyrite physically extracted from the Cismon section
220 sample.

221 **SIMS sulfur isotope analyses**

222 After pre-sputtering by Cs⁺ bombardment for 300 seconds with a 1 nA beam current at the
223 desired raster size, sulfur isotopic ratio experiments were performed in "scanning ion imaging
224 mode" by Cs⁺ bombardment (beam diameter of <1 µm, current of ~10 pA) of raster squares
225 of 20-130 µm, using an electron multiplier (EM) detector on a CAMECA IMS 7f-GEO
226 (Fitchburg, WI, USA) at Washington University in St. Louis to collect counts of ³²S⁻ and ³⁴S⁻
227 for each pixel (grids of 128x128 or 256x256 pixels) for between 20 and 960 planes (1 minute
228 per plane).

229 The size of each exposed grain analyzed was measured using calibrated optical
230 microscope images before analyses. The SIMS stage was x-y calibrated to a stitched optical
231 microscope image of the epoxy puck using digital video camera footage of the gold-coated
232 sample surface in the analysis chamber. Raw isotope ratios for each grain were calculated by
233 taking the mean ³⁴S⁻/³²S⁻ ion count ratio of a central area of the grain over the multiple

234analysis planes. Various corrections were applied to data, including a dead-time correction, an
 235interpolation of $^{34}\text{S}^-$ counts to align in time with those on $^{32}\text{S}^-$, and a quasi-simultaneous
 236arrival (QSA) effect correction.⁵⁷ The magnitude of the QSA undercounting correction is
 237proportional to instrument transmission, i.e., the number of secondary ions reaching the
 238detector per incident primary ion. However, it is not possible to determine primary currents
 239<50 pA very accurately on the 7f-GEO instrument. Therefore, the ratio of the QSA coefficient
 240(β) to the primary ion flux (J) was used to facilitate the correction.⁵⁷ β/J values were
 241determined for each session, via data obtained from the internal standard grains, using the
 242relationship:

$$243 \quad R_{\text{exp}} = R_{\text{cor}} + (\beta/J) \times {}^{34}\text{S}_{\text{exp}} \quad (5)$$

244where R_{exp} and ${}^{34}\text{S}_{\text{exp}}$ are dead time corrected $^{34}\text{S}/^{32}\text{S}$ ratio and ^{34}S count rate, respectively, and
 245 R_{cor} is the QSA corrected $^{34}\text{S}/^{32}\text{S}$ ratio. The instrumental mass fractionation was then corrected
 246for by calculating the mineral-specific fractionation factor ($^{34}\alpha$) based on the mean raw (from
 247SIMS) and expected (from EA-IR-MS) $\delta^{34}\text{S}$ value of the population of internal hydrothermal
 248pyrite or marcasite fragments and dividing the average $^{34}\text{S}/^{32}\text{S}^-$ ratio of each environmental
 249iron sulfide grain by the appropriate $^{34}\alpha$.

250Results

251Extraction procedure

252The grain size distribution of the powdered hydrothermal pyrite crystal ('pre') suggested that
 25392.5% of grains were $>0.5 \mu\text{m}$, the smallest size expected to settle in our setup, so this was
 254set as the expected maximum level of recovery. Recovery of initial pyrite (summarized in
 255Table S1) was highest for 'post' ($58 \pm 12\%$, 1σ) and for 'postSmQz' (57%), but lower for
 256'postQz' ($23 \pm 4\%$, 1σ). Initial extractions ('a') recovered most pyrite, and subsequent
 257extractions ('b' and 'c') were generally not effective in recovering the remaining pyrite. At
 258grain sizes larger than $1.1 \mu\text{m}$ (close to the lower limit of what can be precisely measured on
 259the 7f-GEO; Figure 1), grain size distributions of pre, post, postSmQz and postQz ('1a') were
 260all similar, within the power of our technique to resolve differences (Figure 1, inset).
 261Therefore, the extraction procedure does not impose a grain-size bias for grains $\geq 1.1 \mu\text{m}$.

262[insert Figure 1]

263 Recovery of pyrite for the Santa Barbara Basin sediment sample was estimated to be
 26441.3% by comparing extract mass to previously measured total S abundance.⁵⁰ Extract purity
 265for the Cismon section FeS_2 was estimated using EA to be 61% and by comparing extract
 266mass to CRS abundance, FeS_2 recovery was estimated to be 54.5%, which again closely
 267matches measured recovery for our synthetic sediment samples. In addition, the physically
 268extracted iron sulfides had a bulk $\delta^{34}\text{S}$ value of $-42.5 \pm 0.2\%$, whereas the bulk untreated
 269sample had a $\delta^{34}\text{S}$ value of $-42.1 \pm 0.2\%$.⁵¹ The agreement between the isotopic compositions
 270of the chemically and physically extracted iron sulfides indicates that the physical extraction
 271procedure did not impart any isotopic bias on the population of iron sulfides in the sample.

272SIMS sulfur isotope experiments

273Hydrothermal pyrite

274Sub-angular, randomly oriented 1-3 μm diameter fragments of the hydrothermal pyrite
275(Figure 2A) were analyzed in a 50 μm raster over 375 cycles (30 seconds integration time per
276ion, per cycle; Figure 2B; see Jones et al⁵⁸ for justification). The fragment-to-fragment
277reproducibility ($n=14$, 1σ) was $\pm 1.9\text{‰}$ (Figure 1C; Figure S1A), and using the EA-IRMS
278bulk $\delta^{34}\text{S}$ value of $-1.0 \pm 0.2\text{‰}$, $^{34}\alpha_{\text{pyrite}}$ was calculated to be 0.9964 (Table S2). The standard
279error associated with individual fragments was ± 0.9 to $\pm 3.3\text{‰}$ (1σ ; Figure 2C), with an
280average of $\pm 2.1\text{‰}$ (Figure S1A), and was better for larger fragments due to better counting
281statistics (Figure S2). Lateral intra-fragment reproducibility in the largest fragment (Figure
282S3A) was $\pm 2.8\text{‰}$ ($n=8$, 1σ ; Figure S3B, C), with an average standard error of $\pm 3.0\text{‰}$ (1σ ;
283Figure S3C).

284[insert Figure 2]

285Hydrothermal marcasite

286Sub-angular, randomly oriented 1-5 μm diameter fragments of the hydrothermal marcasite
287crystal (Figure 3A) were analyzed in a 50 μm raster over 375 cycles (30 seconds integration
288time per ion, per cycle; Figure 3B). The fragment-to-fragment reproducibility ($n=25$, 1σ) was
289 $\pm 2.3\text{‰}$ (Figure 3C; Figure S1B), and using the EA-IRMS bulk $\delta^{34}\text{S}$ value of $+4.6 \pm 0.2\text{‰}$,
290 $^{34}\alpha_{\text{marcasite}}$ was calculated to be 1.0014 (Table S2). The standard error associated with
291individual fragments was between ± 1.1 to $\pm 5.2\text{‰}$ (1σ ; Figure 3C), with an average of $\pm 2.6\text{‰}$
292(Figure S1B), and was better for larger fragments due to improved counting statistics (Figure
293S2). Lateral intra-fragment variability in the largest fragment (Figure S4A) was $\pm 1.9\text{‰}$ ($n=6$,
294 1σ ; Figure S4B, C), with an average standard error of $\pm 3.6\text{‰}$ (1σ ; Figure S4C).

295[insert Figure 3]

296Modern sedimentary pyrites (Santa Barbara Basin)

297Optical microscope images (Figure 4A) suggest that the majority of pyrites in this modern
298sediment sample are 1-80 μm -diameter irregular aggregates of intergrown euhedral-to-
299anhedral microcrystals. A minority of pyrites are framboidal (i.e., 5-10 μm -diameter pseudo-
300spheroidal aggregates of equant, equidimensional, non-intergrown microcrystals), while some
301grains display a mixture of the framboidal and irregular textures (Figure 4A). The one
302pristine framboid measured by SIMS (Grain 27; Figure 4B-C) had a $\delta^{34}\text{S}_{\text{pyrite}}$ value of $-42.7 \pm$
303 1.5‰ (1σ ; Figure 4C), the irregular aggregates (Figure 4B) had an average $\delta^{34}\text{S}_{\text{pyrite}}$ value of
304 $+24.2 \pm 2.9\text{‰}$ (1σ , $n=47$; Figure 4C), and five grains displaying mixed textures (Grains 1-2,
30517, 19, and 22; Figure 4B) had $\delta^{34}\text{S}_{\text{pyrite}}$ values between $+0.1 \pm 0.8\text{‰}$ (1σ) and $+16.4 \pm 0.7\text{‰}$
306(1σ ; Figure 4C; Table S2). Within the mixed textured grains, the framboidal areas were
307usually depleted in ^{34}S with respect to the irregular areas (Figure S5). Within the solely
308irregular aggregates, variability in $\delta^{34}\text{S}$ was minimal ($\pm 3.5\text{‰}$, 1σ , $n=11$; Area 14 in Figure 4;
309Figure S6) and comparable to the average standard error associated with measurements
310($\pm 2.6\text{‰}$, 1σ ; Figure S6D). Where present, the apparent variability had no consistent
311directionality (Figure S6). The average standard error associated with individual fragments
312was $\pm 1.5\text{‰}$ (1σ ; Figures 4C, D). The sample average $\delta^{34}\text{S}_{\text{pyrite}}$ value was $+21.7 \pm 10.3\text{‰}$ (1σ ;

313n=53; Figure 4D). The $\delta^{34}\text{S}_{\text{pyrite}}$ variability in this sample (Figures 4C, D) overlaps the bulk
314 $\delta^{34}\text{S}_{\text{CRS}}$ value for the sample of +16.1‰,⁵⁰ and is not normally distributed (Figure 4D).

315[insert Figure 4]

316*Ancient sedimentary pyrites and marcasites (Cismon and Demerara Rise)*

317Optical microscope images of mounted extract from Cismon (Figure S7) display the presence
318of both euhedral marcasite and framboidal pyrite, as supported by laser Raman microprobe
319spot analyses and SEM (Figure S8). Pyrite and marcasite grains from Cismon measured by
320SIMS had average $\delta^{34}\text{S}$ values of $-41.9 \pm 5.2\text{‰}$ (1σ ; n=113) and $-48.5 \pm 5.1\text{‰}$ (1σ ; n=161;
321Figure 5), and cemented pyrite aggregates had an average $\delta^{34}\text{S}$ value of $-42.2 \pm 0.6\text{‰}$ (1σ ;
322n=2), as compared with the bulk $\delta^{34}\text{S}$ composition of -42.1‰ .⁵¹ There was little intra-grain
323 $\delta^{34}\text{S}$ variation in marcasite grains (average $1\sigma = \pm 2.9\text{‰}$, compared to an average intra-grain
324standard error of $\pm 3.5\text{‰}$, 1σ ; Figure S9A-D), and very little intra-grain $\delta^{34}\text{S}$ variation in pyrite
325grains (average $1\sigma = \pm 3.7\text{‰}$, compared to an average intra-grain standard error of $\pm 2.9\text{‰}$, 1σ ;
326Figure S9E-H).

327[insert Figure 5]

328 As with the Cismon section sample, optical microscope images of mounted extract
329from Demerara (Figure S10) display the presence of both euhedral marcasite (in this case
330mostly in irregular aggregates) and framboidal pyrite, as supported by laser Raman
331microprobe spot analyses and SEM (Figure S8). Pyrite and marcasite grains measured by
332SIMS had average $\delta^{34}\text{S}$ values of $-24.9 \pm 11.2\text{‰}$ (1σ ; n=45) and $-26.1 \pm 6.1\text{‰}$ (1σ ; n=19;
333Figure 6), as compared with a bulk $\delta^{34}\text{S}$ composition of $-24.4 \pm 0.2\text{‰}$.⁵² In contrast to those
334from the Cismon section sample, pyrites from the Demerara sample sometimes featured
335larger, more easily-resolved intra-grain variation in $\delta^{34}\text{S}$ (average $1\sigma = \pm 6.0\text{‰}$, compared to
336an average intra-grain standard error of $\pm 1.6\text{‰}$, 1σ ; Figures 6, S11A-D). This was even more
337obviously the case for the marcasites (average $1\sigma = \pm 13.3\text{‰}$, compared to an average intra-
338grain standard error of $\pm 2.1\text{‰}$, 1σ ; Figures 6, S11E-H). Generally, the largest relative
339enrichments in ^{34}S were found near the edges of pyrite framboids (Figure S11A-B), and in the
340extremities of irregular aggregates of both minerals (Figure S11C-H).

341[insert Figure 6]

342Discussion

343*Evaluating the method*

344Despite incomplete recovery (reasons for which are discussed in the Supplement), the
345similarity between artificial pyrite grain size distributions before and after being treated with
346the extraction procedure (Figure 1) suggests that this protocol does not bias pyrite grain size
347distributions, even when grains are at the lower end of sizes that can be measured using a 7f-
348GEO SIMS instrument ($\sim 1 \mu\text{m}$) or when they are mixed with large proportions of quartz
349grains. This is important because where grain size variability exists in populations of iron
350sulfide grains in sedimentary samples, such variability may be coupled to isotopic

351variability;²⁸ thus, a size bias in extraction protocol could produce an isotopic bias in the
352measured results relative to the parent sample.

353 Analysis of fragments of hydrothermal pyrite and marcasite crystals, used as S-
354isotopic standards for our other SIMS experiments, resulted in little inter- (Figures S1-2) or
355intra-fragment (Figures S3-4) variability in $\delta^{34}\text{S}$ (i.e., the standard deviation of measurements
356between or within fragments was always smaller than the average standard error associated
357with those inter- or intra-grain measurements). Therefore, the method is suitable for the
358detection of the potentially large variations in $\delta^{34}\text{S}$ that may exist within or between
359sedimentary iron sulfide grains.^{23,24,28,29,37}

360 The average SIMS $\delta^{34}\text{S}$ values for iron sulfides from samples from the Cismón section
361($-45.8 \pm 6.1\%$, 1σ ; $n=274$; Figure 5) and the Demerara Rise ($-25.3 \pm 11.2\%$, 1σ ; $n=45$;
362Figure 6) are close to the previously reported bulk $\delta^{34}\text{S}_{\text{CRS}}$ values (-42.1% , and -24.4% ,
363respectively).^{51,52} The Santa Barbara Basin pyrites' average $\delta^{34}\text{S}$ value ($+21.7 \pm 10.3\%$, 1σ ;
364 $n=53$; Figure 4) is 5.6% higher than the bulk $\delta^{34}\text{S}_{\text{CRS}}$ value ($+16.1\%$),⁵⁰ but the large range of
365grain-specific values (-42.7% to $+28.9\%$) overlaps with the bulk value, and corresponds to a
366textural dichotomy between isotopically light framboids and isotopically heavy irregular
367aggregates. Accordingly, the discrepancy between bulk and average grain-specific $\delta^{34}\text{S}$ values
368is likely the result of insufficiently representative sampling of the two textural components
369for SIMS analysis. The extraction and analytical procedures introduced in this study are not
370likely to give rise to any isotopic biases, though areas analysed by SIMS may not always be
371entirely representative of the bulk iron sulfides in a sample. In the future, this could be
372rectified by ensuring that SIMS analyses of each texture are in proportion to their relative
373abundances.

374 Importantly, the method documents $\delta^{34}\text{S}$ variability within a sample, which is critical
375for interpreting bulk 'CRS' $\delta^{34}\text{S}$ values. Inter-grain (Figures 4-7), intra-grain (Figures S5-6,
376S9, S11), and inter-mineralogy (Figures 5-6) variability all contribute to a single bulk $\delta^{34}\text{S}$
377value from CRS extraction (which integrates both pyrite and marcasite, as well as potentially
378other reduced S phases). Each type of variability should be considered and investigated when
379making environmental interpretations based on $\delta^{34}\text{S}$ data.

380[insert Figure 7]

381 ***Interpreting sedimentary $\delta^{34}\text{S}$ data***

382 The three samples studied here are distinct in terms of their bulk $\delta^{34}\text{S}$ values and internal $\delta^{34}\text{S}$
383variability (Figure 7). Thus, they can be used to map out the environmental phase space that
384can be explored using this SIMS ion imaging method. The first step to this process is to
385consider that iron sulfide minerals formed in marine sediments record the $\delta^{34}\text{S}$ composition of
386reduced sulfur-bearing aqueous phases (e.g., hydrogen sulfide or polysulfides) in the fluid
387from which they nucleate or grow.^{8,41} In most cases, this fluid is likely to be pore water in
388marine sediments, but pyrites can also precipitate from sulfide-rich water columns.⁵⁹
389Secondly, iron sulfide mineral growth requires a source of iron. Different sources (i.e.,
390mineral or aqueous) are differentially reactive to aqueous reduced sulfur species,⁶⁰ and

391therefore the extent to which iron sulfide minerals record spatio-temporal changes in the $\delta^{34}\text{S}$
392composition of aqueous sulfur is highly dependent on the abundance, mineralogy, and
393speciation of available iron.⁸

394 The sample from Santa Barbara Basin comprises silt-clay sized material from one of
395the “gray layers” that have been hypothesized to represent extremely rapid deposition
396associated with major flood events.^{61–63} Rapid deposition results in pore water sulfate being
397consumed faster than it is supplied.²² We predict a broad range of pyrite $\delta^{34}\text{S}$ values
398corresponding to progressive pyrite precipitation in such a scenario. Berelson et al⁵⁰ found
399that iron disulfides from the sample are highly ^{34}S -enriched in bulk, and are predominantly
400irregular aggregates of pyrite. In this study, we found that the irregular aggregates noted by
401Berelson et al⁵⁰ are even more highly-enriched in ^{34}S than the bulk sample (Figure 4), and
402feature very little intra-grain $\delta^{34}\text{S}$ variability (Figure S6). Conversely, framboidal pyrites are
403depleted in ^{34}S , and feature irregular overgrowths that are usually relatively enriched in ^{34}S
404(Figure S5). This clearly implies a spatial or temporal separation in the growth of these two
405textures. The ^{34}S -depleted framboids likely formed in a relatively open, sulfate-replete
406system, such as pore waters close to the sediment-water interface, or in the water column
407(e.g., in sinking particles harboring anoxic microenvironments⁶⁴). In contrast, the ^{34}S -enriched
408irregular aggregates likely formed in a system in which rapid MSR led to ^{34}S -enriched
409aqueous sulfate and sulfide.^{22,65} MSR must have greatly outpaced pyrite formation, leading to
410the build up of an aqueous reduced sulfur pool in pore fluids with a similar S-isotopic
411composition to the initial sulfate reservoir. The majority of pyrite in the sample precipitated
412from this aqueous reduced sulfur pool. The high rate of MSR relative to the rate of pyrite
413formation was very likely a result of the slow kinetics of iron reduction.⁶⁰

414 Sediments from the Cismom section are thought to represent nearly the opposite end-
415member case to Santa Barbara Basin, with very low bulk CRS $\delta^{34}\text{S}$ values (avg. -42.7%).^{51,52}
416These values have been interpreted to represent pyrite formation in a system that was open to
417the diffusive supply of sulfate,^{66,67} so a narrow intra-sample range of $\delta^{34}\text{S}$ values is expected.
418Accordingly, we find that the sample lacks both inter-grain (Figures 5, 7) and intra-grain
419(Figure S9) $\delta^{34}\text{S}$ variability. Individual pyrite grain $\delta^{34}\text{S}$ values ($-41.9 \pm 5.2\%$, 1σ ; $n=113$)
420and marcasite grain $\delta^{34}\text{S}$ values ($-48.5 \pm 5.1\%$, 1σ ; $n=161$; Figure 5) cluster close to the bulk
421CRS $\delta^{34}\text{S}$ value, suggesting that the S-isotopic composition of aqueous reduced sulfur species
422in the fluid from which the iron sulfides formed was not highly variable. This is consistent
423with a scenario in which all iron sulfides (pyrite framboids, pyrite cements and marcasite
424euhedra) formed either in the water column or in sediments where the rate of sulfate diffusion
425from the water column exceeded the rate of sulfate consumption by MSR.

426 Finally, sediments from the Demerara Rise are thought to represent an intermediate
427case between the Cismom section and Santa Barbara Basin, with a corresponding bulk CRS
428 $\delta^{34}\text{S}$ value of -24.4% for our sample. This black shale sample (422.8 m depth, ODP Leg 207,
429Hole 1258a) was deposited under a euxinic water column, so sulfate consumption by MSR
430very likely outpaced the diffusive replenishment of sulfate.⁵² Therefore, a larger intra-sample
431range of $\delta^{34}\text{S}$ values relative to the Cismom section sample is expected. Accordingly, the

432sample from Demerara Rise features a large amount of $\delta^{34}\text{S}$ variability – both inter- (Figures
4336, 7) and intra-grain (Figure S11), in pyrites and marcasites. The mean $\delta^{34}\text{S}$ values of
434marcasite and pyrite grains (Figure 6) are very similar. However, unlike in the Cismon
435section sample, both pyrite framboids (Figure S11A), and pyrite (Figure S11C) and marcasite
436clusters (Figure S11E, G), have large internal ranges in $\delta^{34}\text{S}$, of $\sim 10\%$ to 65% , with more ^{34}S -
437enriched material on the outer layer of framboids or in discrete zones in the extremities of
438clusters. This pattern is evidence for the formation of these grains over a time interval with
439evolving pore fluid $\delta^{34}\text{S}$. As in the case of the Santa Barbara Basin sample, these results
440suggest that MSR led to ^{34}S -enrichment of aqueous sulfate and sulfide with time in the
441sediment.^{22,65} We also see a ‘tail’ of relatively ^{34}S -enriched pyrites in the $\delta^{34}\text{S}$ distribution
442(Figure 7), which suggests that there must have been sufficiently abundant and reactive iron
443available to sustain iron sulfide mineral growth as pore waters became progressively ^{34}S -
444enriched.^{8,60}

445 Marcasite is thought to be a relatively rare mineral in sedimentary rocks, most notably
446occurring in black shales.^{68,69} The environmental implications of its presence are still poorly
447understood. Experimental work suggests that a very low pH is required for its formation.⁷⁰
448Given that pyrite oxidation can give rise to acidic conditions,⁷¹ this has led to the suggestion
449that the partial oxidation of pre-existing pyrite led to the formation of marcasite in black
450shales.^{68,69} However, given that black shales are thought to represent low $p\text{O}_2$ conditions in
451water column, it is unclear how early diagenetic pyrite oxidation could have occurred in these
452sediments. In addition, the general isotopic similarity between the highly abundant marcasite
453in the Demerara and Cismon samples used in this study, and the coexisting pyrite (Figures 5-
4546), suggests that the two minerals share a formation history. Therefore, marcasite formation
455was probably not a result of low pH conditions created by pyrite oxidation. Another way to
456generate more acidic pore waters than are typical for marine sediments is oxic organic matter
457respiration.⁷¹ This process could have been exacerbated by the high organic loading at both
458sites.⁵² Future work using SIMS will provide further constraints on the relative abundance of
459marcasite and the genetic relationship(s) between pyrite and marcasite in black shales.

460 In conclusion, the methods detailed here provide a basis for the physical extraction
461and accurate and precise sulfur isotopic measurement of microcrystalline iron sulfide grains
462within a range of sediments and sedimentary strata. The minimal size-biasing during the
463extraction procedure suggests that the iron sulfide extracts obtained should be isotopically
464representative of the bulk sample, as confirmed by our samples. The low apparent inter-grain
465variability between isotopically identical pyrite (and marcasite) micro-fragments implies that
466the sulfur isotope composition of diverse environmental populations of $\geq 1\ \mu\text{m}$ -sized pyrites
467and marcasites can be measured accurately and precisely by SIMS ion imaging. Micrometer-
468scale intra-grain variability in $\delta^{34}\text{S}$ can also be measured. Ultimately, it is possible to use this
469approach to unpack local environmental (geochemical and depositional), metabolic, and
470diagenetic signals recorded in iron sulfide grains in both unlithified sediments and rocks.
471With this in mind, the method could be applied to the both the modern and ancient bulk
472sedimentary records of $\delta^{34}\text{S}$ to test previous wide-ranging interpretations of apparent
473excursions in bulk data.⁷²⁻⁷⁴

474 Acknowledgements

475 The authors thank Crystal LaFlamme and one anonymous reviewer, whose insightful
476 comments greatly improved this manuscript. We also thank Itay Halevy, and the members of
477 the Fike and Bradley Labs at Washington University, whose enthusiastic discussions yielded
478 improvements to this work. RNB acknowledges support from the McDonnell Center for the
479 Space Sciences at Washington University in St. Louis. We also acknowledge funding to DAF
480 from DOE/BER (#DE-SC0014613) and from the donors of the American Chemical Society
481 Petroleum Research Fund (#57548-ND2) for partial support of this research.

482 References

4831. LaFlamme C, Martin L, Jeon H, et al. In situ multiple sulfur isotope analysis by SIMS
484 of pyrite, chalcopyrite, pyrrhotite, and pentlandite to refine magmatic ore genetic
485 models. *Chem Geol.* 2016;444:1-15. doi:10.1016/j.chemgeo.2016.09.032.
4862. Newhouse WH. Opaque oxides and sulphides in common igneous rocks. *Bull Geol
487 Soc Am.* 1936;47(1):1-52. doi:10.1130/GSAB-47-1.
4883. Large RR, Maslennikov V V., Robert F, Danyushevsky L V., Chang Z. Multistage
489 sedimentary and metamorphic origin of pyrite and gold in the Giant Sukhoi log
490 deposit, Lena Gold Province, Russia. *Econ Geol.* 2007;102(7):1233-1267.
491 doi:10.2113/gsecongeo.102.7.1233.
4924. Berner RA. Burial of organic carbon and pyrite sulfur in the modern ocean: Its
493 geochemical and environmental significance. *Am J Sci.* 1982;282(4):451-473.
494 doi:10.2475/ajs.282.4.451.
4955. Thode HG, Monster J, Dunford HB. Sulphur isotope geochemistry. *Geochim
496 Cosmochim Acta.* 1961;25(3):159-174. doi:10.1016/0016-7037(61)90074-6.
4976. Berner RA. Modeling atmospheric O₂ over Phanerozoic time. *Geochim Cosmochim
498 Acta.* 2001;65(5):685-694. doi:10.1016/S0016-7037(00)00572-X.
4997. Canfield DE, Farquhar J. Animal evolution, bioturbation, and the sulfate concentration
500 of the oceans. *Proc Natl Acad Sci.* 2009;106(20):8123-8127.
501 doi:10.1073/pnas.0902037106.
5028. Fike DA, Bradley AS, Rose C V. Rethinking the Ancient Sulfur Cycle. *Annu Rev Earth
503 Planet Sci.* 2015;43(1):593-622. doi:10.1146/annurev-earth-060313-054802.
5049. Ding T, Bai R, Li Y, Wan D, Zou X, Zhang Q. Determination of the absolute ³²S/³⁴S
505 ratio of IAEA-S-1 reference material and V-CDT sulfur isotope standard. *Sci China,
506 Ser D Earth Sci.* 1999;42(1):45-51. doi:10.1007/BF02878497.
50710. Canfield DE, Raiswell R, Westrich JT, Reaves CM, Berner RA. The use of chromium
508 reduction in the analysis of reduced inorganic sulfur in sediments and shales. *Chem
509 Geol.* 1986;54(1-2):149-155. doi:10.1016/0009-2541(86)90078-1.
51011. Fossing H, Jørgensen BB. Measurement of bacterial sulfate reduction in sediments:
511 Evaluation of a single-step chromium reduction method. *Biogeochemistry.*
512 1989;8(3):205-222. doi:10.1007/BF00002889.
51312. Giesemann A, Jäger HJ, Norman AL, Krouse HR, Brand WA. On-Line Sulfur-Isotope
514 Determination Using an Elemental Analyzer Coupled to a Mass Spectrometer. *Anal
515 Chem.* 1994;66(18):2816-2819. doi:10.1021/ac00090a005.

51613. Morrison J, Fourel F, Churchman D. Isotopic sulphur analysis by CF-IRMS. 2000.
517 Available at: [https://scholar.google.com/scholar?](https://scholar.google.com/scholar?hl=en&as_sdt=0%2C26&q=Morrison+Isotopic+sulphur+analysis+by+CF-IRMS.&btnG=)
518 [hl=en&as_sdt=0%2C26&q=Morrison+Isotopic+sulphur+analysis+by+CF-](https://scholar.google.com/scholar?hl=en&as_sdt=0%2C26&q=Morrison+Isotopic+sulphur+analysis+by+CF-IRMS.&btnG=)
519 [IRMS.&btnG=](https://scholar.google.com/scholar?hl=en&as_sdt=0%2C26&q=Morrison+Isotopic+sulphur+analysis+by+CF-IRMS.&btnG=). Accessed July 4, 2018.
52014. Grassineau N V., Matthey DP. Measurement of sulphur isotopic compositions of
521 sulphide minerals using new continuous He-flow EA-MS technology. In: *Goldschmidt*
522 *Conference.*; 1998:537-538. Available at:
523 http://rruff.info/doclib/MinMag/Volume_62A/62A-1-537.pdf. Accessed July 4, 2018.
52415. Grassineau N V., Matthey DP, Lowry D. Sulfur isotope analysis of sulfide and sulfate
525 minerals by continuous flow-isotope ratio mass spectrometry. *Anal Chem.*
526 2001;73(2):220-225. doi:10.1021/ac000550f.
52716. Grassineau N V. High-precision EA-IRMS analysis of S and C isotopes in geological
528 materials. *Appl Geochemistry.* 2006;21(5):756-765.
529 doi:10.1016/j.apgeochem.2006.02.015.
53017. Studley SA, Ripley EM, Elswick ER, et al. Analysis of sulfides in whole rock matrices
531 by elemental analyzer- continuous flow isotope ratio mass spectrometry. *Chem Geol.*
532 2002;192(1-2):141-148. doi:10.1016/S0009-2541(02)00162-6.
53318. Clough R, Evans P, Catterick T, Evans EH. $\delta^{34}\text{S}$ measurements of sulfur by
534 multicollector inductively coupled plasma mass spectrometry. *Anal Chem.*
535 2006;78(17):6126-6132. doi:10.1021/ac060875h.
53619. Mason PRD, Košler J, de Hoog JCM, Sylvester PJ, Meffan-Main S. In situ
537 determination of sulfur isotopes in sulfur-rich materials by laser ablation multiple-
538 collector inductively coupled plasma mass spectrometry (LA-MC-ICP-MS). *J Anal At*
539 *Spectrom.* 2006;21(2):177-186. doi:10.1039/B510883G.
54020. Craddock PR, Rouxel OJ, Ball LA, Bach W. Sulfur isotope measurement of sulfate and
541 sulfide by high-resolution MC-ICP-MS. *Chem Geol.* 2008;253(3-4):102-113.
542 doi:10.1016/j.chemgeo.2008.04.017.
54321. Paris G, Sessions AL, Subhas A V., Adkins JF. MC-ICP-MS measurement of $\delta^{34}\text{S}$ and
544 $\Delta^{33}\text{S}$ in small amounts of dissolved sulfate. *Chem Geol.* 2013;345:50-61.
545 doi:10.1016/j.chemgeo.2013.02.022.
54622. Claypool GE. Ventilation of marine sediments indicated by depth profiles of pore
547 water sulfate and $\delta^{34}\text{S}$. *Geochemical Soc Spec Publ.* 2004;9(C):59-65.
548 doi:10.1016/S1873-9881(04)80007-5.
54923. Fischer WW, Fike DA, Johnson JE, et al. SQUID-SIMS is a useful approach to
550 uncover primary signals in the Archean sulfur cycle. *Proc Natl Acad Sci.*
551 2014;111(15):5468-5473. doi:10.1073/pnas.1322577111.
55224. Marin-Carbonne J, Remusat L, Sforza MC, Thomazo C, Cartigny P, Philippot P. Sulfur
553 isotope's signal of nanopyrites enclosed in 2.7 Ga stromatolitic organic remains reveal
554 microbial sulfate reduction. *Geobiology.* 2018;16(2):121-138. doi:10.1111/gbi.12275.
55525. Johnston DT, Farquhar J, Canfield DE. Sulfur isotope insights into microbial sulfate
556 reduction: When microbes meet models. *Geochim Cosmochim Acta.*
557 2007;71(16):3929-3947. doi:10.1016/j.gca.2007.05.008.
55826. Leavitt WD, Halevy I, Bradley AS, Johnston DT. Influence of sulfate reduction rates
559 on the Phanerozoic sulfur isotope record. *Proc Natl Acad Sci.* 2013;110(28):11244-
560 11249. doi:10.1073/pnas.1218874110.

56127. Jorgensen BB. A theoretical model of the stable sulfur isotope distribution in marine
562 sediments. *Geochim Cosmochim Acta*. 1979;43(3):363-374. doi:10.1016/0016-
563 7037(79)90201-1.
56428. Gomes ML, Fike DA, Bergmann KD, Jones C, Knoll AH. Environmental insights
565 from high-resolution (SIMS) sulfur isotope analyses of sulfides in Proterozoic
566 microbialites with diverse mat textures. *Geobiology*. 2018;16(1):17-34.
567 doi:10.1111/gbi.12265.
56829. Drake H, Whitehouse MJ, Heim C, et al. Unprecedented³⁴S-enrichment of pyrite
569 formed following microbial sulfate reduction in fractured crystalline rocks.
570 *Geobiology*. 2018;16(5):556-574. doi:10.1111/gbi.12297.
57130. Cui H, Brussel VU, Kitajima K, Fournelle J, Ishida A. Questioning the Biogenicity of
572 Neoproterozoic Superheavy Pyrite by SIMS. *Am Mineral*. 2018;103:1362-1400.
573 doi:10.2138/am-2018-6489.
57431. Riciputi LR, Paterson BA, Ripperdan RL. Measurement of light stable isotope ratios
575 by SIMS: *Int J Mass Spectrom*. 1998;178(1-2):81-112. doi:10.1016/S1387-
576 3806(98)14088-5.
57732. Greenwood JP, Mojzsis SJ, Coath CD. Sulfur isotopic compositions of individual
578 sulfides in Martian meteorites ALH840001 and Nakhla: Implications for crust-regolith
579 exchanges on Mars. *Earth Planet Sci Lett*. 2000;184(1):23-35. doi:10.1016/S0012-
580 821X(00)00301-0.
58133. Papineau D, Mojzsis SJ, Coath CD, Karhu JA, McKeegan KD. Multiple sulfur
582 isotopes of sulfides from sediments in the aftermath of Paleoproterozoic glaciations.
583 *Geochim Cosmochim Acta*. 2005;69(21):5033-5060. doi:10.1016/j.gca.2005.07.005.
58434. Whitehouse MJ, Kamber BS, Fedo CM, Lepland A. Integrated Pb- and S-isotope
585 investigation of sulphide minerals from the early Archaean of southwest Greenland.
586 *Chem Geol*. 2005;222(1-2):112-131. doi:10.1016/j.chemgeo.2005.06.004.
58735. Kamber BS, Whitehouse MJ. Micro-scale sulphur isotope evidence for sulphur cycling
588 in the late Archean shallow ocean. *Geobiology*. 2007;5(1):5-17. doi:10.1111/j.1472-
589 4669.2006.00091.x.
59036. Kozdon R, Kita NT, Huberty JM, Fournelle JH, Johnson CA, Valley JW. In situ sulfur
591 isotope analysis of sulfide minerals by SIMS: Precision and accuracy, with application
592 to thermometry of ~3.5Ga Pilbara cherts. *Chem Geol*. 2010;275(3-4):243-253.
593 doi:10.1016/j.chemgeo.2010.05.015.
59437. Drake H, Åström ME, Tullborg EL, Whitehouse M, Fallick AE. Variability of sulphur
595 isotope ratios in pyrite and dissolved sulphate in granitoid fractures down to 1km depth
596 - Evidence for widespread activity of sulphur reducing bacteria. *Geochim Cosmochim*
597 *Acta*. 2013;102:143-161. doi:10.1016/j.gca.2012.10.036.
59838. Meyer NR, Zerkle AL, Fike DA. Sulphur cycling in a Neoarchaean microbial mat.
599 *Geobiology*. 2017;15(3):353-365. doi:10.1111/gbi.12227.
60039. Fu J, Hu Z, Li J, et al. Accurate determination of sulfur isotopes ($\delta^{33}\text{S}$ and $\delta^{34}\text{S}$) in
601 sulfides and elemental sulfur by femtosecond laser ablation MC-ICP-MS with non-
602 matrix matched calibration. *J Anal At Spectrom*. 2017;32(12). doi:10.1039/c7ja00282c.
60340. Zhu ZY, Jiang SY, Ciobanu CL, Yang T, Cook NJ. Sulfur isotope fractionation in pyrite
604 during laser ablation: Implications for laser ablation multiple collector inductively
605 coupled plasma mass spectrometry mapping. *Chem Geol*. 2017;450:223-234.

- 606 doi:10.1016/j.chemgeo.2016.12.037.
60741. Rickard D. *Sulfidic Sediments and Sedimentary Rocks*. Elsevier B.V.; 2012.
608 doi:10.1016/B978-0-444-52989-3.00017-9.
60942. Valley JW, Kita NT. In-situ oxygen isotope geochemistry by Ion Microprobe. *Mineral
610 Assoc Canada Short Course*. 2009;41(May):19-63. Available at:
611 <https://pdfs.semanticscholar.org/44d4/b868830727ad1ed366b5dcb95789ca4994fa.pdf>.
612 Accessed September 25, 2018.
61343. Flinter BH. The magnetic separation of some alluvial minerals in Malaya. *Am Mineral*.
614 1959;44:738-751. Available at: [https://pubs.geoscienceworld.org/msa/ammin/article-
615 abstract/44/7-8/738/541523](https://pubs.geoscienceworld.org/msa/ammin/article-abstract/44/7-8/738/541523). Accessed August 3, 2018.
61644. Vallentyne JR. Isolation of pyrite spherules from recent sediments. *Limnol Oceanogr*.
617 1963;8(1):16-30. doi:10.4319/lo.1963.8.1.0016.
61845. Volkov II, Fomina LS. Influence of organic material and processes of sulfide formation
619 on distribution of some trace elements in deep-water sediments of Black Sea. In: *The
620 Black Sea - Geology, Chemistry and Biology*.; 1974:456-476. Available at:
621 [http://archives.datapages.com/data/specpubs/sedimen1/data/a145/a145/0001/0450/045
622 6.htm](http://archives.datapages.com/data/specpubs/sedimen1/data/a145/a145/0001/0450/0456.htm). Accessed August 3, 2018.
62346. Raiswell R, Plant J. The incorporation of trace elements into pyrite during diagenesis
624 of black shales, Yorkshire, England. *Econ Geol*. 1980;75(5):684-699.
625 doi:10.2113/gsecongeo.75.5.684.
62647. Ivor Roberts F. Trace element chemistry of pyrite: A useful guide to the occurrence of
627 sulfide base metal mineralization. *J Geochemical Explor*. 1982;17(1):49-62.
628 doi:10.1016/0375-6742(82)90019-X.
62948. Proske U, Wood R, Fallon S, Stevenson J. Use of heavy liquid density separation to
630 remove pyrite from sediment samples for radiocarbon dating. *Quat Geochronol*.
631 2015;25:66-71. doi:10.1016/j.quageo.2014.10.002.
63249. Imai H, Kawasaki M, Yamaguchi M, Takahashi M. Mineralization and paragenesis of
633 the Huanzala mine, central Peru. *Econ Geol*. 1985;80(2):461-478.
634 doi:10.2113/gsecongeo.80.2.461.
63550. Berelson WM, Morine L, Sessions AL, Rollins N, Fleming JC, Schwalbach J. Santa
636 Barbara Basin Flood Layers: Impact on Sediment Diagenesis. *SEPM Spec Publ No
637 110*. 2018. doi:10.2110/sepmsp.110.11.
63851. Gomes ML, Hurtgen MT, Sageman BB. Biogeochemical sulfur cycling during
639 Cretaceous oceanic anoxic events: A comparison of OAE1a and OAE2.
640 *Paleoceanography*. 2016;31(2):233-251. doi:10.1002/2015PA002869.
64152. Raven MR, Fike DA, Bradley AS, Gomes ML, Owens JD, Webb SL. Paired organic
642 matter and pyrite $\delta^{34}\text{S}$ records reveal mechanisms of carbon, sulfur, and iron cycle
643 disruption during Ocean Anoxic Event 2. doi:10.31223/OSF.IO/K97BW.
64453. Schneider CA, Rasband WS, Eliceiri KW. NIH Image to ImageJ: 25 years of image
645 analysis. *Nat Methods*. 2012;9(7):671-675. doi:10.1038/nmeth.2089.
64654. Loveday J. Methods for analysis of irrigated soils. *Commonw Bur Soils B2 -
647 Commonw Bur Soils*. 1973:38-62. Available at: [http://agris.fao.org/agris-
648 search/search.do?recordID=XE7528376%0Ahttps://ezp.lib.unimelb.edu.au/login?
649 url=https://search.ebscohost.com/login.aspx?
650 direct=true&db=edsref&AN=CBS.LOVEDAY.COMMONWEALTHAGRICULTURA](http://agris.fao.org/agris-search/search.do?recordID=XE7528376%0Ahttps://ezp.lib.unimelb.edu.au/login?url=https://search.ebscohost.com/login.aspx?direct=true&db=edsref&AN=CBS.LOVEDAY.COMMONWEALTHAGRICULTURA)

- 651 LB.AIGC.CH&site=eds-live. Accessed August 3, 2018.
65255. Lentfer CJ, Cotter MM, Boyd WE. Particle settling times for gravity sedimentation
653 and centrifugation: A practical guide for Palynologists. *J Archaeol Sci.*
654 2003;30(2):149-168. doi:10.1006/jasc.2001.0786.
65556. Bryant RN, Pasteris JD, Fike DA. Variability in the Raman Spectrum of Unpolished
656 Growth and Fracture Surfaces of Pyrite Due to Laser Heating and Crystal Orientation.
657 *Appl Spectrosc.* 2018;72(1):37-47. doi:10.1177/0003702817736516.
65857. Jones C, Fike DA, Peres P. Investigation of the quasi-simultaneous arrival (QSA)
659 effect on a CAMECA IMS 7f-GEO. *Rapid Commun Mass Spectrom.* 2017;31(7):623-
660 630. doi:10.1002/rcm.7828.
66158. Jones C, Fike DA, Meyer KM. SIMS Methodology for Isotopic Ratio Measurement of
662 Micro-Grains in Thin Sections: True Grain Size Estimation and Deconvolution of
663 Inter-Grain Size Gradients and Intra-Grain Radial Gradients. *Geostand Geoanalytical*
664 *Res.* 2018. doi:10.1111/ggr.12247.
66559. Suits NS, Wilkin RT. Pyrite formation in the water column and sediments of a
666 meromictic lake. *Geology.* 1998;26(12):1099-1102. doi:10.1130/0091-
667 7613(1998)026<1099:PFITWC>2.3.CO;2.
66860. Meyers SR. Production and preservation of organic matter: The significance of iron.
669 *Paleoceanography.* 2007;22:PA4211. doi:10.1029/2006PA001332.
67061. Fleischer P. Mineralogy and sedimentation history, Santa Barbara Basin, California. *J*
671 *Sediment Res.* 1972;42(1):49-58. doi:10.1306/74D72487-2B21-11D7-
672 8648000102C1865D.
67362. Thornton SE. Origin of mass flow sedimentary structures in hemipelagic basin
674 deposits: Santa Barbara Basin, California Borderland. *Geo-Marine Lett.* 1986;6(1):15-
675 19. doi:10.1007/BF02311691.
67663. Li C, Sessions AL, Valentine DL, Thiagarajan N. Organic Geochemistry D / H
677 variation in terrestrial lipids from Santa Barbara Basin over the past 1400 years : A
678 preliminary assessment of paleoclimatic relevance. *Org Geochem.* 2011;42(1):15-24.
679 doi:10.1016/j.orggeochem.2010.09.011.
68064. Bianchi D, Weber TS, Kiko R, Deutsch C. Global niche of marine anaerobic
681 metabolisms expanded by particle microenvironments. *Nat Geosci.* 2018;11(4):263-
682 268. doi:10.1038/s41561-018-0081-0.
68365. Aller RC, Heilbrun C, Panzeca C, Zhu Z, Baltzer F. Coupling between sedimentary
684 dynamics, early diagenetic processes, and biogeochemical cycling in the Amazon-
685 Guianas mobile mud belt: Coastal French Guiana. In: *Marine Geology.* Vol 208.;
686 2004:331-360. doi:10.1016/j.margeo.2004.04.027.
68766. Bellanca A, Claps M, Erba E, et al. Orbitally induced limestone/marlstone rhythms in
688 the albian-cenomanian cismon section (Venetian region, northern Italy):
689 Sedimentology, calcareous and siliceous plankton distribution, elemental and isotope
690 geochemistry. *Palaeogeogr Palaeoclimatol Palaeoecol.* 1996;126(3-4):227-260.
691 doi:10.1016/S0031-0182(96)00041-7.
69267. Gambacorta G, Bersezio R, Weissert H, Erba E. Onset and demise of Cretaceous
693 oceanic anoxic events: The coupling of surface and bottom oceanic processes in two
694 pelagic basins of the western Tethys. *Paleoceanography.* 2016;31(6):732-757.
695 doi:10.1002/2015PA002922.

69668. Schieber J. Oxidation of detrital pyrite as a cause for Marcasite Formation in marine
697 lag deposits from the Devonian of the eastern US. *Deep Res Part II Top Stud*
698 *Oceanogr.* 2007;54(11-13):1312-1326. doi:10.1016/j.dsr2.2007.04.005.
69969. Schieber J. Marcasite in Black Shales--a Mineral Proxy for Oxygenated Bottom
700 Waters and Intermittent Oxidation of Carbonaceous Muds. *J Sediment Res.*
701 2011;81(7):447-458. doi:10.2110/jsr.2011.41.
70270. Murowchick JB, Barnes HL. Effects of temperature and degree of supersaturation on
703 pyrite morphology. *Am Mineral.* 1987;72:24-1250. Available at:
704 http://www.minsocam.org/ammin/AM72/AM72_1241.pdf. Accessed March 12, 2016.
70571. Soetaert K, Hofmann AF, Middelburg JJ, Meysman FJR, Greenwood J. Reprint of
706 "The effect of biogeochemical processes on pH." *Mar Chem.* 2007;106(1-2):380-401.
707 doi:10.1016/j.marchem.2007.06.008.
70872. Parnell J, Boyce AJ, Mark D, Bowden S, Spinks S. Early oxygenation of the terrestrial
709 environment during the Mesoproterozoic. *Nature.* 2010;468(7321):290-293.
710 doi:10.1038/nature09538.
71173. Gill BC, Lyons TW, Young SA, Kump LR, Knoll AH, Saltzman MR. Geochemical
712 evidence for widespread euxinia in the Later Cambrian ocean. *Nature.*
713 2011;469(7328):80-83. doi:10.1038/nature09700.
71474. Pasquier V, Sansjofre P, Rabineau M, Revillon S, Houghton J, Fike DA. Pyrite sulfur
715 isotopes reveal glacial–interglacial environmental changes. *Proc Natl Acad Sci.*
716 2017;114(23):5941-5945. doi:10.1073/pnas.1618245114.
- 717